

# Selenium Supplementation and the Effects on Reproductive Outcomes, Biomarkers of Inflammation, and Oxidative Stress in Women with Polycystic Ovary Syndrome

**Authors** M. Razavi<sup>1</sup>, M. Jamilian<sup>2</sup>, Z. Fakhrieh Kashan<sup>3</sup>, Z. Heidar<sup>4</sup>, M. Mohseni<sup>3</sup>, Y. Ghandi<sup>3</sup>, T. Bagherian<sup>3</sup>, Z. Asemi<sup>3</sup>

**Affiliations** Affiliation addresses are listed at the end of the article

## Key words

- selenium
- supplementation
- polycystic ovary syndrome
- reproductive outcomes
- inflammation
- oxidative stress

## Abstract

Selenium supplementation could be effective on reproductive outcomes, biomarkers of inflammation, and oxidative stress among women with polycystic ovary syndrome (PCOS). The aim of the study was to determine the effects of selenium supplementation on reproductive outcomes, biomarkers of inflammation, and oxidative stress in PCOS patients. The present randomized double-blind, placebo-controlled trial was conducted on 64 women aged 18–40 years old with PCOS at the clinic affiliated to Ardabil University of Medical Sciences, Ardabil, Iran. The participants were randomly assigned to 2 groups receiving either 200 µg selenium daily (n=32) or placebo (n=32) for 8 weeks. Hormonal profiles, biomarkers of inflammation, and oxidative stress were measured and compared both before and after the treatment. After 8 weeks of intervention, pregnancy rate in the selenium

group was higher than in the placebo group: 18.8 (6/32) vs. 3.1% (1/32), p=0.04. In addition, alopecia (40.6 vs. 9.4%, p=0.004) and acne (46.9 vs. 12.5 %, p=0.003) decreased following the consumption of selenium supplements compared with placebo. Additionally, patients who received selenium supplements had significantly decreased serum dehydroepiandrosterone (DHEA) levels (p=0.02), hirsutism (modified Ferriman–Gallwey scores) (p<0.001), serum high sensitivity C-reactive protein (hs-CRP) (p=0.02), and plasma malondialdehyde (MDA) levels (p=0.01) compared with placebo. We did not observe any significant effects of taking selenium supplements on other hormonal profiles, nitric oxide (NO), and other biomarkers of oxidative stress. Taken together, selenium supplementation for 8 weeks among PCOS women had beneficial effects on reproductive outcomes, DHEA, hs-CRP, and MDA levels.

**Supporting Information** for this article is available online at <http://www.thieme-connect.de/products>.

received 16.04.2015  
accepted 02.07.2015

## Bibliography

**DOI** <http://dx.doi.org/10.1055/s-0035-1559604>  
Published online:  
August 12, 2015  
Horm Metab Res 2016;  
48: 185–190  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0018-5043

## Correspondence

**Z. Asemi, PhD**  
Department of Nutrition  
Kashan University of Medical  
Sciences  
Kashan  
I. R. Iran  
Tel.: +98/31/5546 3378  
Fax: +98/31/5546 3377  
asemi\_r@yahoo.com

## Abbreviations

β-HCG	β-Human chorionic gonadotropin
CVD	Cardiovascular disease
DHEA	Dehydroepiandrosterone
FRAP	Ferric reducing antioxidant power
FSH	Follicular-stimulating hormone
FOH	Functional ovarian hyperandrogenism
GSH	Glutathione
GSH-Px	Glutathione peroxidase
Hs-CRP	High-sensitivity C-reactive protein
ITT	Intention-to-treat
LH	Luteinizing hormone
LOCF	Last Observation Carried Forward
MDA	Malondialdehyde
MUFA	Monounsaturated fatty acid
NO	Nitric oxide
OCPs	Oral contraceptives
PCOS	Polycystic ovary syndrome
PUFA	Polyunsaturated fatty acid

ROS	Reactive oxygen species
RDA	Recommended Dietary Allowances
RA	Rheumatoid arthritis
SIRS	Systemic inflammatory response syndrome
SFA	Saturated fatty acid
TAC	Total antioxidant capacity
TAS	Total antioxidant status
T2DM	Type 2 diabetes mellitus
TNF-α	Tumor necrosis factor alpha
TDF	Total dietary fiber
tT	Total testosterone

## Introduction

Polycystic ovary syndrome (PCOS) is a common endocrine and metabolic aberration [1], affecting 5–10 % of women of childbearing age in worldwide [2]. Previous studies have shown that PCOS

is the most prevalent cause of female infertility, which is characterized by ovarian hyperandrogenism and chronic anovulation [3]. In addition, due to increased production of free radicals and reactive oxygen species (ROS) [4], elevated circulating protein carbonyls, tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), and decreased insulin signaling [5,6], it is associated with increased biomarkers of inflammation and oxidative stress. Increased inflammatory factors and biomarkers of oxidative stress among patients with PCOS may predict the development of cardiovascular disease (CVD) and type 2 diabetes mellitus (T2DM) [7–9].

Recent studies have reported that the use of insulin-sensitizing drugs including metformin, rosiglitazone, and pioglitazone improves ovulation by improving clinical and biochemical features of patients diagnosed with PCOS [10]. Selenium is an essential trace element, which has importance to human biology and health. More recently, evidence has also been presented that selenium may affect several reproductive and obstetric complications including male and female infertility [11]. Furthermore, current data supports the beneficial effect of selenium supplementation on metabolic profiles and biomarkers of oxidative stress among patients with PCOS [12, 13]. Valenta et al. [14] demonstrated that high-dose selenium supplementation for 14 days (1000  $\mu$ g on day 1 and 500  $\mu$ g/day on days 2–14) decreased plasma CRP levels in patients with systemic inflammatory response syndrome (SIRS)/sepsis. Increased erythrocyte and plasma total antioxidant status (TAS), reduced glutathione (GSH), and glutathione peroxidase (GSH-Px) were also seen following the intake selenium supplements in patients with epilepsy and refractory epilepsy [15]. However, no significant change in malondialdehyde (MDA) concentrations was seen after the consumption of selenium supplements in rats [16]. Beneficial effects of selenium supplementation on improved reproductive outcomes, biomarkers of inflammation, and oxidative stress might be attributed to its inhibitory effect on augmentation of proinflammatory cytokines and reactive oxygen species/reactive nitrogen species [17,18]. Borderline selenium status is high among healthy Iranian women. Rafraf et al. [19] reported that 69 women have serum selenium concentrations under 80  $\mu$ g/l. Although there is no specific data on mean daily intake of selenium in Iranian women with PCOS, the mean of selenium consumption among postmenopausal women was significantly less than the Recommended Dietary Allowances (RDA) [20]. Considering these points, we hypothesized that selenium supplementation might affect reproductive outcomes, biomarkers of inflammation, and oxidative stress in patients with PCOS. We are aware of no such study among PCOS women. This study was thus aimed to investigate the effect of selenium supplementation on reproductive outcomes, biomarkers of inflammation, and oxidative stress of PCOS women.

## Subjects and Methods

### Participants

Sixty four women with PCOS from October 2014 to December 2014 were included in the patients attending the outpatient clinic of the Infertility Research Center of Ardabil Medical University. The current study was approved by the Institutional Review Board at Ardabil University of Medical Sciences (AUMS). Informed written consent was obtained from all participants. This study was done based on the guidelines laid down in the Declaration of Helsinki. The trial was registered in the Iranian

website ([www.irct.ir](http://www.irct.ir)) for registration of clinical trials (IRCT code: IRCT201412295623N33). The diagnosis of PCOS was according to European Society for Human Reproduction and Embryology/American Society for Reproductive Medicine guidelines (Rotterdam criteria, 2003), as including at least 2 of the following 3 criteria: 1) oligo- and/or anovulation as the presence of chronic amenorrhea or a menstrual cycle length of less than 21 days or more than 35 days, or more than 4 days variation between cycles; 2) clinical or biochemical signs of hyperandrogenism; and 3) polycystic ovary morphology shown on ultrasound examination, defined as 12 or more small follicles [21]. Our inclusion criteria were age between 18 and 40 years with PCOS according to Rotterdam criteria. In the current study, we excluded patients who had elevated levels of prolactin, thyroid disorder, or T2DM and congenital adrenal hyperplasia. In addition, all PCOS women had normal baseline renal function tests, bilirubin, and aminotransferases. During the last 3 months before the intervention, none of the study patients used any form of oral contraceptives (OCs), other steroid hormones, or any other treatments likely to affect ovarian function, insulin sensitivity, or inflammatory factors, biomarkers of oxidative stress as well as nutritional supplements.

### Study design

At the first trial and after stratification for pre-supplementation BMI (<25 and  $\geq$  25 kg/m<sup>2</sup>) and age (<30 and  $\geq$  30 years), patients with PCOS were randomly divided into 2 groups. Group 1 received 200  $\mu$ g daily selenium tablet as selenium yeast (n=32) and group 2 received the placebo (n=32) for 8 weeks. Selenium supplements and its placebos (cellulose) were manufactured by Nature Made Pharmaceutical Company (California, USA) and Barij Essence Pharmaceutical Company (Kashan, Iran), respectively. Supplements and placebos were in the same form of package and the patients and researcher were not conscious of the content of the pack until the end of trial. In addition, all patients with PCOS took metformin at the initial dose of 500 mg, which was elevated in a stepwise manner during the first 3 weeks to incorporate the side effects until the patients were taking a total of 1500 mg/day. Patient allocation and block size were obtained using random number tables. At the time of randomization, sequentially numbered, sealed envelopes were opened. Allocation to study group was concealed until the main analyses were completed. The randomized allocation sequence, enrolling participants, and assigning participants to interventions were done by a trained midwife at gynecology clinic. At the beginning study, the subjects were requested to keep their usual diet and level of physical activity throughout the study period as well as not to receive any anti-inflammatory medications, supplements, and other medications that might affect their reproductive physiology during the 8-week intervention. The use of selenium supplements and placebos throughout the study was checked through asking participants to bring the medication containers. To increase the compliance, all patients were receiving short messages on their cell phones to take the supplements each day. All participants provided 3 dietary records (1 weekend day and 2 weekdays) and 3 physical activity records at week 2, 4, and 6 of intervention to make sure that they maintained their usual diet and physical activity during intervention. The dietary records were based on estimated values in household measurements. In the current study, we used Nutritionist IV software (First Databank, San Bruno, CA, USA) adjusted for Iranian foods to take macro and micronutrients intakes of patients according to 3-day food diaries.

### Assessment of variables

All participants were evaluated at baseline study on the third day of a spontaneous or progesterone-induced menstrual cycle. Anthropometric assessments included determinations of height, weight (Seca, Hamburg, Germany), and body mass index (BMI). Clinical assessments included determinations of hirsutism using modified Ferriman-Gallwey scores [22], of acne score [23], and of alopecia based on assessment guidelines collated by Olsen et al. [24]. Acne was marked by a 4-point scale: 0, no acne; 1, minor acne on face; 2, moderate acne on face only; and 3, severe acne, face and back or chest. In case of missed menstrual period, we checked  $\beta$ -human chorionic gonadotropin ( $\beta$ -HCG) to detect pregnancy a week after the missed period. If  $\beta$ -HCG levels were higher than 25 (by the Vidas method), the patient was pregnant and medications discontinued. Fasting blood samples (10 ml) were collected before and 8 weeks after of the intervention at Ardabil reference laboratory in an early morning after an overnight fast. Blood was collected in 2 separate tubes: 1) one without EDTA to separate the serum, in order to quantify serum prolactin, follicular-stimulating hormone (FSH) and luteinizing hormone (LH), free testosterone, dehydroepiandrosterone (DHEA) and high sensitivity C-reactive protein (hs-CRP) levels and 2) another one containing EDTA to examine plasma nitric oxide (NO) and biomarkers of oxidative stress. Blood samples were immediately centrifuged (Hettich, 78532 Tuttlingen, Germany) at 3 500 rpm for 10 min to separate the serum. The samples were then stored at  $-70^{\circ}\text{C}$  until being analyzed at the AUMS reference laboratory. Commercial kits were used to measure serum prolactin, FSH and LH concentrations (Pars Azmun, Tehran, Iran). All inter- and intra-assay CVs for prolactin, FSH and LH measurements were less than 7%. Serum free testosterone, DHEA, and 17-OH progesterone concentrations were determined by using commercial kits (Monobind, CA, USA). Serum hs-CRP was quantified using an ELISA kit (LDN, Nordhorn, Germany) with intra- and interassay CVs of 2.9 and 4.8%, respectively. The plasma NO concentration was determined by the Griess method [25]. Plasma total antioxidant capacity (TAC) was assessed by the use of the ferric reducing antioxidant power (FRAP) method developed by Benzie and Strain [26]. Plasma GSH was examined using the method of Beutler et al. [27]. The plasma MDA levels were determined by the thiobarbituric acid reactive substance spectrophotometric test [28]. CVs for plasma TAC, GSH and MDA were 1.1, 2.8, and 3.7%, respectively.

### Statistical methods

We used the Kolmogorov-Smirnov test to examine if variables were normally distributed. Intention-to-treat (ITT) analysis of the primary study end-point was performed for all the randomly assigned participants. Missing data from dropped out participants were imputed using the method of "Last Observation Carried Forward (LOCF)". The Pearson chi-square test was used to compare categorical variables. Independent sample Student's *t*-test was used to detect differences in general characteristics and dietary intake between the 2 groups. To determine the effects of selenium supplementation on hormonal profiles, inflammatory factors and biomarkers of oxidative stress, one-way repeated-measures ANOVA was used to evaluate the between-group changes in variables during the study. In this analysis, the treatment was regarded as between-subject factor and time with 2 time-points (baseline and 8 weeks after of the intervention) was considered as within-subject factor. To assess if the magnitude of the change in dependent variables depended

on the baseline age and BMI, we controlled all analyses for baseline values of age and BMI to avoid potential bias. These analyses were also done using one-way repeated measures ANOVA. A *p*-value  $<0.05$  was considered statistically significant. All statistical analyses were done using the Statistical Package for Social Science version 17 (SPSS Inc., Chicago, IL, USA). In the current study, the primary endpoint variables were biomarkers of oxidative stress including plasma TAC, GSH, and MDA levels. To calculate the sample size, we used the standard formula suggested for parallel clinical trials by considering type one error ( $\alpha$ ) of 0.05 and type 2 error ( $\beta$ ) of 0.20 (power=80%). Based on a previous study [29], we used  $2.19\mu\text{mol/l}$  as SD and  $2.80\mu\text{mol/l}$  as the difference in mean (*d*) of MDA levels as key variable. Based on this, we reached to 27 patients in each group. Assuming a dropout of 5 patients per group, the final sample size was determined to be 32 patients per group.

## Results



### Baseline evaluations and dropouts

In the current study, 64 women with PCOS met the inclusion criteria based on Rotterdam criteria and were enrolled in the study. Among the individuals in the selenium group, 2 women [withdrawn due to personal reasons ( $n=2$ )] and in the placebo group, 2 women [withdrawn due to personal reasons ( $n=2$ )] did not complete the trial. However, as the analysis was done according to ITT protocol, all 64 patients with PCOS were included in the end analysis. On the average, the rate of compliance in our study was high, such that  $>90\%$  of tablets were taken throughout the study in both groups. No side effects were reported following the administration of selenium supplements in patients with PCOS throughout the study.

### General characteristics of study participants

Mean age ( $25.1\pm 4.5$  vs.  $25.4\pm 4.9$  years,  $p=0.85$ ), baseline BMI ( $24.7\pm 3.5$  vs.  $25.3\pm 4.3\text{ kg/m}^2$ ,  $p=0.53$ ), and end-of-trial BMI ( $24.3\pm 3.5$  vs.  $25.1\pm 4.2\text{ kg/m}^2$ ,  $p=0.47$ ) were not significantly different between selenium and placebo groups.

### Dietary intakes of study participants throughout the study

According to the 3-day dietary records taken throughout the intervention, any significant change was not seen between the 2 groups in terms of dietary intakes of energy, carbohydrates, proteins, fats, saturated fatty acids (SFAs), polyunsaturated fatty acids (PUFAs), monounsaturated fatty acids (MUFAs), cholesterol, total dietary fiber (TDF), selenium, magnesium, manganese, vitamins C and E (data not shown).

### Clinical outcomes

After 8 weeks of intervention, pregnancy rate in the selenium group was higher than in the placebo group: 18.8 (6/32) vs. 3.1% (1/32),  $p=0.04$  (Table S1). In addition, alopecia (40.6 vs. 9.4%,  $p=0.004$ ) and acne (46.9 vs. 12.5%,  $p=0.003$ ) decreased following the consumption of selenium supplements compared with placebo.

### The effect of selenium supplementations on biochemical measurements

Patients who received selenium supplements had significantly decreased serum DHEA levels ( $p=0.02$ ), hirsutism ( $p<0.001$ ),

serum hs-CRP ( $p=0.02$ ), and plasma MDA levels ( $p=0.01$ ) compared with placebo (Table 1). We did not observe any significant effects of taking selenium supplements on other hormonal profiles, NO, and other biomarkers of oxidative stress. We controlled the baseline levels in the analyses. However, after adjustment no significant changes in our findings occurred, except for FSH ( $p=0.03$ ) and DHEA levels ( $p=0.12$ ).

### Adjusted changes in biochemical measurements

Additional adjustments for age and baseline BMI did not affect our findings, except for FSH ( $p=0.03$ ) and DHEA levels ( $p=0.13$ ) (Table S2).

A summary of patient flow diagram is presented in Fig. S1.

## Discussion

We revealed for the first time that selenium supplementation in PCOS women resulted in a significant increase in pregnancy rate, a decreased reduction in alopecia, acne, DHEA, hirsutism, hs-CRP and MDA levels. It must be considered that mean dietary plus supplemental selenium intake was lower in our study patients than upper limits (400  $\mu\text{g}$ ). However, data on the effects of selenium supplementation on toxicity/teratogenicity even in subjects with high dietary selenium intake are conflicting. For instance, some of the subjects in Burk et al study [30] ingested >800  $\mu\text{g}$  selenium/d for 16 weeks. However, this is considerably more than the Institute of Medicine's tolerable upper level of 400  $\mu\text{g}/\text{d}$  [31], no signs of selenium toxicity in mentioned study were observed. In another study by Reid et al. [32] more subjects on 3200  $\mu\text{g}/\text{day}$  have reported symptoms of selenium toxicity.

Women with PCOS are susceptible to infertility, hirsutism, increased inflammatory factors and increased biomarkers of oxidative stress [33]. Our study demonstrated that taking selenium supplements for 8 weeks in women with PCOS resulted in a significant increase in pregnancy rate, a significant reduction in alopecia and acne compared with placebo. Data regarding favorable effects of selenium intake and female fertility are

scarce. In line with our study, conception rate following the intake of 50 mg/ml selenium as barium selenate was higher than in the control ewes [34]. In addition, fertility has been improved by supplemental co-administration of vitamin E and selenium in cattle [35]. Paszkowski et al. [36] reported that patients with unexplained infertility had a significant decrease in follicular selenium concentrations compared with patients with tubal infertility. Furthermore, a recent study has shown lower levels of serum and follicular fluid selenium in women undergoing in vitro fertilization treatment compared with nonpregnant control women [37]. Edassery et al. [38] reported that women with unexplained infertility or premature ovarian failure have significantly increased serum levels of the ovarian autoantibody protein, selenium binding protein-1. However, some researchers have not found association between herd selenium concentrations and fertility parameters [39]. Previous studies have reported that selenium intake is associated with decreased biomarkers of oxidative stress and insulin resistance in women with PCOS [12,13]. Antioxidants are compounds that are capable of disposing or suppressing the formation of ROS, free radicals, and lipid hydroperoxides [40]. However, no study has directly evaluated the effects of oxidative stress on fertility in women; a recent review has reported that antioxidants may influence female reproduction [41]. In addition, insulin resistance in patients with PCOS influences the developmental potential of human immature oocytes, as indicated by impaired oocyte maturation, fewer fertilized oocytes, and cleaved embryos [42,43].

Results of the present study show that selenium supplementation among PCOS women decreased serum DHEA levels and hirsutism, while did not affect other hormonal profiles. As this study is among the first that reports the effect of selenium supplementation on hormonal profiles among women with PCOS, we did not compare findings of the current study with others although few studies assessed the association between selenium levels and hormonal profiles in women with PCOS. For instance, Coskun et al. [44] reported decreased plasma concentrations of selenium and a negative association between selenium and LH, total testosterone (tT) in PCOS women, indicating that selenium may play a role in the pathogenesis of PCOS related with hyper-

**Table 1** The effect of selenium supplementations on hormonal status, biomarkers inflammation and oxidative stress.

	Placebo group (n=32)			Selenium group (n=32)			p*
	Week 0	Week 8	Change	Week 0	Week 8	Change	
Prolactin (mIU/l)	643.27±376.31	546.11±262.97	-97.15±443.26	697.01±614.53	743.28±585.41	46.26±337.43	0.15
FSH (IU/l)	7.92±2.85	8.40±3.05	0.48±2.55	6.39±3.17	6.15±3.38	-0.24±4.28	0.41
LH (IU/l)	13.67±13.53	11.48±7.72	-2.19±13.77	10.93±8.21	10.60±8.94	-0.33±10.21	0.54
Free testosterone (pg/ml)	3.24±1.77	3.02±1.66	-0.21±0.92	2.70±1.51	2.41±1.46	-0.28±1.15	0.78
DHEA ( $\mu\text{g}/\text{ml}$ )	1.59±0.72	1.56±0.73	-0.02±0.41	2.01±0.80	1.65±0.85 <sup>†</sup>	-0.36±0.73	0.02
17-OH Progesterone (ng/ml)	2.00±1.39	1.76±1.28	-0.23±1.68	2.39±2.01	1.92±1.82	-0.46±2.56	0.66
mF-G scores	8.34±6.30	8.12±6.24	-0.21±0.83	12.46±7.21	10.12±5.45 <sup>†</sup>	-2.34±2.57	<0.001
hs-CRP (ng/ml)	2572.26±2506.08	2765.81±2372.19	193.54±1117.45	2184.06±2693.84	1472.70±1444.43 <sup>†</sup>	-711.35±1959.37	0.02
NO ( $\mu\text{mol}/\text{l}$ )	57.15±17.09	57.22±16.98	0.07±23.38	46.91±7.59	50.76±14.75	3.85±13.37	0.43
TAC (mmol/l)	696.45±137.43	692.13±138.74	-4.32±200.88	823.14±195.00	803.60±147.74	-19.54±135.00	0.72
GSH ( $\mu\text{mol}/\text{l}$ )	437.30±107.12	472.35±148.66	35.05±168.55	476.64±122.77	459.92±120.73	-16.72±132.53	0.17
MDA ( $\mu\text{mol}/\text{l}$ )	4.40±2.71	5.8±3.03 <sup>†</sup>	1.40±3.11	4.95±1.25	4.81±1.36	-0.13±1.03	0.01

All values are means  $\pm$  SDs

\* Obtained from repeated measures ANOVA test (time  $\times$  group interaction)

DHEA: Dehydroepiandrosterone; FSH: Follicle-stimulating hormone; GSH: Glutathione; hs-CRP: High sensitivity C-reactive protein; LH: Luteinizing hormone; mF-G: Modified Ferriman-Gallwey; MDA: Malondialdehyde; NO: Nitric oxide; TAC: Total antioxidant capacity

<sup>†</sup> Significant difference with baseline study

androgenism. Selenium supplementation may result in decreased serum DHEA levels and hirsutism through improved markers of insulin metabolism [13] and decreased oxidative stress [12]. Previous studies have demonstrated that a positive association between hyperandrogenism, hyperinsulinemia, and insulin resistance in females with elevated androgen levels due to PCOS [45,46]. In addition, oxidative stress is directly correlated with both insulin resistance and hyperandrogenism, which in turn contribute to endocrine and biochemical alterations in women with functional ovarian hyperandrogenism (FOH) [47]. The present study revealed that taking selenium supplements in PCOS patients for 8 weeks was associated with decreased serum hs-CRP levels, but did not affect plasma NO concentrations. Supporting with our study, administration of selen plus supplements contained 50µg selenium, 8 mg zinc, 400µg vitamin A, 125 mg vitamin C, and 40 mg vitamin E daily has reduced hs-CRP concentrations among rheumatoid arthritis (RA) for 12 weeks [48]. In addition, high-dose selenium supplementation for 14 days (1000µg on day 1 and 500µg/day on days 2–14) has led to a significant decrease in plasma CRP levels in patients with SIRS/sepsis [14]. Unlike, no significant effect on hs-CRP levels was observed after intake of cereal biscuit with selenized onion, curcuma, and green tea for 2 months among healthy adults [49]. Similar finding was also seen in centrally obese women who consumed 200µg selenium supplements for 6 weeks [50]. Increased inflammatory factors in patients with PCOS render them at a potential increased risk for the development of atherosclerosis, T2DM, cancer, infertility, and other comorbidities [51]. Selenium intake may decrease serum hs-CRP levels through inhibiting the activation of NF-kappa B by modulating selenoprotein genes expression [52] and increasing selenoprotein biosynthesis which in turn results in suppressed CRP production [53]. We found that selenium supplementation among women with PCOS has resulted in a significant reduction in plasma MDA levels, but it could not affect plasma TAC and GSH levels. Similarly, Salehi et al. [54] demonstrated a significant decrease in MDA levels after administration of 200µg selenium supplements per day for 12 weeks among hemodialysis patients. Furthermore, vitamin E-plus-selenium combination decreased MDA concentrations in streptozotocin-induced diabetic rats [55]. Supplementation with 200µg selenium per day for 3 weeks did not influence biomarkers of oxidative stress including TAS and GSH levels among overweight adults [56]. However, few researchers did not observe such effects of selenium supplementation on MDA levels. For instance, any significant change in MDA levels was not observed after taking selenium supplements in rats [16]. Previous studies have shown that oxidative stress in female reproduction is associated with polycystic ovarian syndrome and endometriosis [57]. Needless to say, these pathologies negatively influence pregnancy rates and IVF outcomes. Selenium is an essential component of the erythrocyte GSH-Px system [58], which functions as part of an antioxidant defense to protect polyunsaturated fatty acids, the damaging effects of free radicals, and lipid hydroperoxides such as MDA [59]. In addition, selenium might decrease hydroperoxides levels via inhibiting production of proinflammatory cytokines and reactive oxygen species/reactive nitrogen species [17]. Taken together, selenium supplementation for 8 weeks among PCOS women had beneficial effects on reproductive outcomes, DHEA, hs-CRP and MDA levels.

## Author Contributions

Z.A. contributed in conception, design, statistical analysis, and drafting of the manuscript. M.R, M.J., Z.F., Z.H., M.M., Y.Gh. and T.B. contributed in data collection and manuscript drafting.

## Acknowledgements

The study was supported by a grant (No. 9306) from Ardabil University of Medical Sciences.

## Conflict of Interest

None of the authors had any personal or financial conflict of interest.

## Affiliations

- <sup>1</sup> Department of Gynecology and Obstetrics, School of Medicine, Ardabil University of Medical Sciences, Ardabil, I. R. Iran
- <sup>2</sup> Department of Gynecology and Obstetrics, School of Medicine, Arak University of Medical Sciences, Arak, I. R. Iran
- <sup>3</sup> Research Center for Biochemistry and Nutrition in Metabolic Diseases, Kashan University of Medical Sciences, Kashan, I. R. Iran
- <sup>4</sup> Department of Gynecology and Obstetrics, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, I. R. Iran

## References

- 1 Asemi Z, Esmailzadeh A. DASH diet, insulin resistance, and serum hs-CRP in polycystic ovary syndrome: a randomized controlled clinical trial. *Horm Metab Res* 2015; 47: 232–238
- 2 Homburg R. Polycystic ovary syndrome. *Best Pract Res Clin Obstet Gynaecol* 2008; 22: 261–274
- 3 Amooee S, Parsanezhad ME, Ravanbod Shirazi M, Alborzi S, Samsami A. Metformin versus chromium picolinate in clomiphene citrate-resistant patients with PCOs: A double-blind randomized clinical trial. *Iran J Reprod Med* 2013; 11: 611–618
- 4 Macut D, Bjekic-Macut J, Savic-Radojevic A. Dyslipidemia and oxidative stress in PCOS. *Front Horm Res* 2013; 40: 51–63
- 5 Gonzalez F, Rote NS, Minium J, Kirwan JP. Reactive oxygen species-induced oxidative stress in the development of insulin resistance and hyperandrogenism in polycystic ovary syndrome. *J Clin Endocrinol Metab* 2006; 91: 336–340
- 6 Li S, Tao T, Wang L, Mao X, Zheng J, Zhao A, Liu W. The expression of 11beta-HSDs, GR, and H6PDH in subcutaneous adipose tissue from polycystic ovary syndrome subjects. *Horm Metab Res* 2013; 45: 802–807
- 7 Cicchitano P, Dentamaro I, Carbonara R, Bulzisi G, Dachille A, Caputo P, Riccardi R, Locorotondo M, Mandurino C, Matteo Ciccone M. Cardiovascular Risk in Women With PCOS. *Int J Endocrinol Metab* 2012; 10: 611–618
- 8 Caglar GS, Oztas E, Karadag D, Pabuccu R, Demirtas S. Ischemia-modified albumin and cardiovascular risk markers in polycystic ovary syndrome with or without insulin resistance. *Fertil Steril* 2011; 95: 310–313
- 9 Anagnostis P, Efsthathiadou ZA, Gougoura S, Polyzos SA, Karathanasi E, Dritsa P, Kita M, Koukoulis GN. Oxidative stress and reduced antioxidative status, along with endothelial dysfunction in acromegaly. *Horm Metab Res* 2013; 45: 314–318
- 10 Tang T, Lord JM, Norman RJ, Yasmin E, Balen AH. Insulin-sensitising drugs (metformin, rosiglitazone, pioglitazone, D-chiro-inositol) for women with polycystic ovary syndrome, oligo amenorrhoea and subfertility. *Cochrane Database Syst Rev* 2012; 5: CD003053 doi:10.1002/14651858
- 11 Mistry HD, Broughton Pipkin F, Redman CW, Poston L. Selenium in reproductive health. *Am J Obstet Gynecol* 2012; 206: 21–30
- 12 Kose SA, Naziroglu M. Selenium reduces oxidative stress and calcium entry through TRPV1 channels in the neutrophils of patients with polycystic ovary syndrome. *Biol Trace Elem Res* 2014; 158: 136–142
- 13 Jamilian M, Razavi M, Fakhrie Kashan Z, Ghandi Y, Bagherian T, Asemi Z. Metabolic response to selenium supplementation in women with polycystic ovary syndrome: a randomized, double-blind, placebo-controlled trial. *Clin Endocrinol (Oxf)* 2015; 82: 885–891

- 14 Valenta J, Brodska H, Drabek T, Hendl J, Kazda A. High-dose selenium substitution in sepsis: a prospective randomized clinical trial. *Intensive Care Med* 2011; 37: 808–815
- 15 Yurekli VA, Naziroglu M. Selenium and topiramate attenuates blood oxidative toxicity in patients with epilepsy: a clinical pilot study. *Biol Trace Elem Res* 2013; 152: 180–186
- 16 Kang BP, Bansal MP, Mehta U. Selenium supplementation and diet induced hypercholesterolemia in the rat: changes in lipid levels, malondialdehyde production and the nitric oxide synthase activity. *Gen Physiol Biophys* 1998; 17: 71–78
- 17 Zeng J, Zhou J, Huang K. Effect of selenium on pancreatic proinflammatory cytokines in streptozotocin-induced diabetic mice. *J Nutr Biochem* 2009; 20: 530–536
- 18 Ruder EH, Hartman TJ, Goldman MB. Impact of oxidative stress on female fertility. *Curr Opin Obstet Gynecol* 2009; 21: 219–222
- 19 Rafraf M, Mahdavi R, Rashidi MR. Serum selenium levels in healthy women in Tabriz, Iran. *Food Nutr Bull* 2008; 29: 83–86
- 20 Mansour A, Ahadi Z, Qorbani M, Hosseini S. Association between dietary intake and seasonal variations in postmenopausal women. *J Diabetes Metab Disord* 2014; 13: 52
- 21 Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril* 2004; 81: 19–25
- 22 Hatch R, Rosenfield RL, Kim MH, Tredway D. Hirsutism: implications, etiology, and management. *Am J Obstet Gynecol* 1981; 140: 815–830
- 23 Kolodziejczyk B, Duleba AJ, Spaczynski RZ, Pawelczyk L. Metformin therapy decreases hyperandrogenism and hyperinsulinemia in women with polycystic ovary syndrome. *Fertil Steril* 2000; 73: 1149–1154
- 24 Olsen EA, Hordinsky MK, Price VH, Roberts JL, Shapiro J, Canfield D, Duvic M, King Jr LE, McMichael AJ, Randall VA, Turner ML, Sperling L, Whiting DA, Norris D. Alopecia areata investigational assessment guidelines – Part II. National Alopecia Areata Foundation. *J Am Acad Dermatol* 2004; 51: 440–447
- 25 Tatsch E, Bochi GV, Pereira Rda S, Kober H, Agertt VA, de Campos MM, Gomes P, Duarte MM, Moresco RN. A simple and inexpensive automated technique for measurement of serum nitrite/nitrate. *Clin Biochem* 2011; 44: 348–350
- 26 Benzie IF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: the FRAP assay. *Anal Biochem* 1996; 239: 70–76
- 27 Beutler E, Gelbart T. Plasma glutathione in health and in patients with malignant disease. *J Lab Clin Med* 1985; 105: 581–584
- 28 Janero DR. Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury. *Free Radic Biol Med* 1990; 9: 515–540
- 29 Asemi Z, Jamilian M, Mesdaghinia E, Esmailzadeh A. Effects of selenium supplementation on glucose homeostasis, inflammation and oxidative stress in gestational diabetes: randomized, double-blind, placebo-controlled trial. *Nutrition* 2015; DOI: <http://dx.doi.org/10.1016/j.nut.2015.04.014>
- 30 Burk RF, Norworthy BK, Hill KE, Motley AK, Byrne DW. Effects of chemical form of selenium on plasma biomarkers in a high-dose human supplementation trial. *Cancer Epidemiol Biomarkers Prev* 2006; 15: 804–810
- 31 Johnson LJ, Meacham SL, Kruskall LJ. The antioxidants – vitamin C, vitamin E, selenium, and carotenoids. *J Agromedicine* 2003; 9: 65–82
- 32 Reid ME, Stratton MS, Lillico AJ, Fakih M, Natarajan R, Clark LC, Marshall JR. A report of high-dose selenium supplementation: response and toxicities. *J Trace Elem Med Biol* 2004; 18: 69–74
- 33 Bargiota A, Diamanti-Kandarakis E. The effects of old, new and emerging medicines on metabolic aberrations in PCOS. *Ther Adv Endocrinol Metab* 2012; 3: 27–47
- 34 Munoz C, Carson AF, McCoy MA, Dawson LE, Irwin D, Gordon AW, Kilpatrick DJ. Effect of supplementation with barium selenate on the fertility, prolificacy and lambing performance of hill sheep. *Vet Rec* 2009; 164: 265–271
- 35 Arechiga CF, Vazquez-Flores S, Ortiz O, Hernandez-Ceron J, Porras A, McDowell LR, Hansen PJ. Effect of injection of beta-carotene or vitamin E and selenium on fertility of lactating dairy cows. *Theriogenology* 1998; 50: 65–76
- 36 Paszkowski T, Traub AI, Robinson SY, McMaster D. Selenium dependent glutathione peroxidase activity in human follicular fluid. *Clin Chim Acta* 1995; 236: 173–180
- 37 Ozkaya MO, Naziroglu M, Barak C, Berkkanoglu M. Effects of multivitamin/mineral supplementation on trace element levels in serum and follicular fluid of women undergoing in vitro fertilization (IVF). *Biol Trace Elem Res* 2011; 139: 1–9
- 38 Edassery SL, Shatavi SV, Kunkel JP, Hauer C, Brucker C, Penumatsa K, Yu Y, Dias JA, Luborsky JL. Autoantigens in ovarian autoimmunity associated with unexplained infertility and premature ovarian failure. *Fertil Steril* 2010; 94: 2636–2641
- 39 Jukola E, Hakkainen J, Saloniemi H, Sankari S. Blood selenium, vitamin E, vitamin A, and beta-carotene concentrations and udder health, fertility treatments, and fertility. *J Dairy Sci* 1996; 79: 838–845
- 40 Ruder EH, Hartman TJ, Blumberg J, Goldman MB. Oxidative stress and antioxidants: exposure and impact on female fertility. *Hum Reprod Update* 2008; 14: 345–357
- 41 Ebisch IM, Thomas CM, Peters WH, Braat DD, Steegers-Theunissen RP. The importance of folate, zinc and antioxidants in the pathogenesis and prevention of subfertility. *Hum Reprod Update* 2007; 13: 163–174
- 42 Kovacs G, Wood C. The current status of polycystic ovary syndrome. *Aust N Z J Obstet Gynaecol* 2001; 41: 65–68
- 43 Vlaisavljevic V, Kovac V, Sajko MC. Impact of insulin resistance on the developmental potential of immature oocytes retrieved from human chorionic gonadotropin-primed women with polycystic ovary syndrome undergoing in vitro maturation. *Fertil Steril* 2009; 91: 957–959
- 44 Coskun A, Arikani T, Kilinc M, Arikani DC, Ekerbicer HC. Plasma selenium levels in Turkish women with polycystic ovary syndrome. *Eur J Obstet Gynecol Reprod Biol* 2013; 168: 183–186
- 45 Hauner H, Ditschuneit HH, Pal SB, Moncayo R, Pfeiffer EF. Fat distribution, endocrine and metabolic profile in obese women with and without hirsutism. *Metabolism* 1988; 37: 281–286
- 46 Muntzer J, Hofer D, Trummer C, Ulbing M, Harger A, Pieber T, Owen L, Keevil B, Brabant G, Lerchbaum E, Obermayer-Pietsch B. Testosterone to dihydrotestosterone ratio as a new biomarker for an adverse metabolic phenotype in the polycystic ovary syndrome. *J Clin Endocrinol Metab* 2015; 100: 653–660
- 47 Suresh S, Vijayakumar T. Correlations of Insulin Resistance and Serum Testosterone Levels with LH:FSH Ratio and Oxidative Stress in Women with Functional Ovarian Hyperandrogenism. *Indian J Clin Biochem* 2015; 30: 345–350
- 48 Jalili M, Kolahi S, Aref-Hosseini SR, Mamegani ME, Hekmatdoost A. Beneficial role of antioxidants on clinical outcomes and erythrocyte antioxidant parameters in rheumatoid arthritis patients. *Int J Prev Med* 2014; 5: 835–840
- 49 Madaric A, Kadrabova J, Krajcovicova-Kudlackova M, Valachovicova M, Spustova V, Mislanova C, Kajaba I, Blazicek P. The effect of bioactive complex of quercetin, selenium, catechins and curcumin on cardiovascular risk markers in healthy population after a two month consumption. *Bratislav Lek Listy* 2013; 114: 84–87
- 50 Alizadeh M, Safaeiyan A, Ostadrahimi A, Estakhri R, Daneghian S, Ghafari A, Gargari BP. Effect of L-arginine and selenium added to a hypocaloric diet enriched with legumes on cardiovascular disease risk factors in women with central obesity: a randomized, double-blind, placebo-controlled trial. *Ann Nutr Metab* 2012; 60: 157–168
- 51 Pawelczak M, Rosenthal J, Milla S, Liu YH, Shah B. Evaluation of the Pro-inflammatory Cytokine Tumor Necrosis Factor-alpha in Adolescents with Polycystic Ovary Syndrome. *J Pediatr Adolesc Gynecol* 2014; 27: 356–359
- 52 He YT, Liu DW, Ding LY, Li Q, Xiao YH. Therapeutic effects and molecular mechanisms of anti-fibrosis herbs and selenium on rats with hepatic fibrosis. *World J Gastroenterol* 2004; 10: 703–706
- 53 Duntas LH. Selenium and inflammation: underlying anti-inflammatory mechanisms. *Horm Metab Res* 2009; 41: 443–447
- 54 Salehi M, Sohrobi Z, Ekramzadeh M, Fallahzadeh MK, Ayatollahi M, Geramizadeh B, Hassanzadeh J, Sagheb MM. Selenium supplementation improves the nutritional status of hemodialysis patients: a randomized, double-blind, placebo-controlled trial. *Nephrol Dial Transplant* 2013; 28: 716–723
- 55 Guney M. Selenium-vitamin E combination modulates endometrial lipid peroxidation and antioxidant enzymes in streptozotocin-induced diabetic rat. *Biol Trace Elem Res* 2012; 149: 234–240
- 56 Savory LA, Kerr CJ, Whiting P, Finer N, McEneny J, Ashton T. Selenium supplementation and exercise: effect on oxidant stress in overweight adults. *Obesity (Silver Spring)* 2012; 20: 794–801
- 57 Gupta S, Ghulmiyyah J, Sharma R, Halabi J, Agarwal A. Power of proteomics in linking oxidative stress and female infertility. *Biomed Res Int* 2014; 9: 16212
- 58 Ozturk IC, Batcioglu K, Karatas F, Hazneci E, Genc M. Comparison of plasma malondialdehyde, glutathione, glutathione peroxidase, hydroxyproline and selenium levels in patients with vitiligo and healthy controls. *Indian J Dermatol* 2008; 53: 106–110
- 59 Rohr-Udilova N, Sieghart W, Eferl R, Stoiber D, Bjorkhem-Bergman L, Eriksson LC, Stolze K, Hayden H, Keppler B, Sagmeister S, Grasl-Kraupp B, Schulte-Hermann R, Peck-Radosavljevic M. Antagonistic effects of selenium and lipid peroxides on growth control in early hepatocellular carcinoma. *Hepatology* 2012; 55: 1112–1121